

Forward Jets and Forward W Production at Hadron Colliders

Jeppe R. Andersen

Institute for Particle Physics Phenomenology
University of Durham, UK

in collaboration with V. Del Duca, S. Frixione, F. Maltoni, C. Schmidt, and
W.J. Stirling

Outline of the talk

- Short introduction to BFKL at Hadron Colliders: Leading Logs and its Problems with Momentum Conservation

Outline of the talk

- Short introduction to BFKL at Hadron Colliders: Leading Logs and its Problems with Momentum Conservation
- Construction of a BFKL MC
Momentum conserved, running coupling effects, experimental cuts

Outline of the talk

- Short introduction to BFKL at Hadron Colliders: Leading Logs and its Problems with Momentum Conservation
- Construction of a BFKL MC
Momentum conserved, running coupling effects, experimental cuts
- Application of the MC (and fixed order QCD) to a dijet study: dijet and $W + 2$ jet production at large rapidity separation

Why BFKL at Hadron Colliders


BFKL (**B**alitskii, **F**adin, **K**uraev, **L**ipatov):
resummation of large logarithms in the
perturbation series for processes with two large
and disparate energy scales

BFKL most known from the study of the structure
function F_2 in DIS


However, in the study of the structure function F_2 ,
nothing guarantees that the logarithms resummed
in the BFKL approach are kinematically favoured

A better study of BFKL effects is forward jet
production in DIS or dijet production at hadron
colliders (as suggested by Mueller and Navelet)

BFKL at Hadron Colliders



High energy
limit: $\frac{\hat{s}}{|\hat{t}|} \rightarrow \infty$



BFKL resums to all orders terms in the perturbative expansion of the form


$$\left(\alpha_s \ln \frac{\hat{s}}{|\hat{t}|} \right)^n \sim (\alpha_s \Delta y)^n$$

BFKL at LL

gluon-gluon scattering:

$$\frac{d\hat{\sigma}_{gg}}{d^2\vec{p}_{T_a} d^2\vec{p}_{T_b}} = \underbrace{\left[\frac{C_A \alpha_s}{p_{T_a}^2} \right]}_{\text{QCD IF}} \underbrace{f(\vec{q}_a, \vec{q}_b, \Delta y)}_{\text{BFKL effects}} \underbrace{\left[\frac{C_A \alpha_s}{p_{T_b}^2} \right]}_{\text{QCD IF}}$$

$$\vec{q}_a = \vec{p}_{T_a}, \vec{q}_b = -\vec{p}_{T_b}$$



Resum leading logarithms in $\frac{\hat{s}}{|\hat{t}|} \sim \Delta y$ contributing to $f(\vec{q}_a, \vec{q}_b, \Delta y)$.

Take a good look at f !

The BFKL Equation

Mellin transformed in Δy

$$\tilde{f}(\vec{q}_a, \vec{q}_b, \omega) = \int_0^\infty d\Delta y e^{-\omega\Delta y} f(\vec{q}_a, \vec{q}_b, \Delta y)$$

$$\omega \tilde{f}(\vec{q}_a, \vec{q}_b, \omega) = \frac{1}{2} \delta(\vec{q}_a - \vec{q}_b) + \frac{C_A \alpha_s}{\pi^2} \int \frac{d^2 \vec{k}}{\vec{k}^2} K(\vec{q}_a, \vec{q}_b, \vec{k}),$$


where the kernel K is given by

$$K(\vec{q}_a, \vec{q}_b, \vec{k}) = \tilde{f}(\vec{q}_a + \vec{k}, \vec{q}_b, \omega) - \frac{\vec{q}_a^2}{\vec{k}^2 + (\vec{q}_a + \vec{k})^2} \tilde{f}(\vec{q}_a, \vec{q}_b, \omega).$$

IR divergences from **real** and **virtual** gluon radiation cancel.

The BFKL Equation

Solve the BFKL equation **analytically** by **integrating** over the **full \vec{k} phase space** for gluon emission and allowing **any number** of gluons to radiate.



$$\hat{\sigma}_{gg} = \frac{\pi C_A^2 \alpha_s^2}{2P_{T,\min}^2} \frac{e^{\lambda \Delta y}}{\sqrt{\pi B \Delta y}}, \quad B = 14\zeta(3)\bar{\alpha}_s, \quad \lambda = \frac{\alpha_s C_A}{\pi} 4 \ln 2 \approx 0.45$$

Energy & Mom. conservation

Problem of including pdf's to evaluate hadronic cross section. So far we have $\frac{d\hat{\sigma}}{d\Delta y}$, need $\frac{d\sigma}{d\Delta y}$, where

$$\sigma = \int dx_a dx_b f(x_a) f(x_b) \hat{\sigma}$$
$$x_a = \frac{P_{T_a}}{\sqrt{s}} e^{y_a} + \frac{P_{T_b}}{\sqrt{s}} e^{y_b} + \sum_{i=1}^n \frac{k_{i\perp}}{\sqrt{s}} e^{y_i}$$
$$x_b = \frac{P_{T_a}}{\sqrt{s}} e^{-y_a} + \frac{P_{T_b}}{\sqrt{s}} e^{-y_b} + \sum_{i=1}^n \frac{k_{i\perp}}{\sqrt{s}} e^{-y_i}$$

These **terms** are inaccessible in the standard BFKL approach:

BFKL gluons are emitted at no cost in energy!

...not just energy conservation

Look for observables that are independent of the PDF's, e.g. **angular decorrelation** of jets.

Need better understanding of event topology.

Experimental cuts on BFKL gluon radiation?

Wishes for a MC:

1. Energy/momentum conservation
2. Handle on gluon radiation (cuts, gaps,...)

It will turn out we can also handle effects from the running of the coupling

(which scale should the coupling be evaluated at in the BFKL ladder?)

Building a BFKL MC

$$\omega \tilde{f}(\vec{q}_a, \vec{q}_b, \omega) = \frac{1}{2} \delta(\vec{q}_a - \vec{q}_b) + \frac{\bar{\alpha}_s}{\pi} \int \frac{d^2 \vec{k}}{\vec{k}^2} K(\vec{q}_a, \vec{q}_b, \vec{k}),$$

$$K(\vec{q}_a, \vec{q}_b, \vec{k}) = \tilde{f}(\vec{q}_a + \vec{k}, \vec{q}_b, \omega) - \frac{\vec{q}_a^2}{\vec{k}^2 + (\vec{q}_a + \vec{k})^2} \tilde{f}(\vec{q}_a, \vec{q}_b, \omega).$$

Want to solve by iteration to “unfold” the gluon contribution. Need to maintain the cancellation of IR div. Introduce “resolution scale”

$\mu \ll P_{T,\min}$. Rewrite the BFKL equation

$$\omega \tilde{f}(\vec{q}_{a\perp}, \vec{q}_{b\perp}, \omega) = \frac{1}{2} \delta(\vec{q}_{a\perp} - \vec{q}_{b\perp}) + \frac{\bar{\alpha}_s}{\pi} \int_{k_{\perp}^2 > \mu^2} \frac{d^2 \vec{k}_{\perp}}{k_{\perp}^2} \tilde{f}(\vec{q}_{a\perp} + \vec{k}_{\perp}, \vec{q}_{b\perp}, \omega)$$

$$+ \frac{\bar{\alpha}_s}{\pi} \int \frac{d^2 \vec{k}_{\perp}}{k_{\perp}^2} \left[\tilde{f}(\vec{q}_{a\perp} + \vec{k}_{\perp}, \vec{q}_{b\perp}, \omega) \theta(\mu^2 - k_{\perp}^2) - \frac{q_{a\perp}^2 \tilde{f}(\vec{q}_{a\perp}, \vec{q}_{b\perp}, \omega)}{k_{\perp}^2 + (\vec{q}_{a\perp} + \vec{k}_{\perp})^2} \right]$$

μ independent!

Since in the last integral $k_{\perp}^2 \leq \mu^2 \ll q_{a\perp}^2, q_{b\perp}^2$ we will neglect \vec{k}_{\perp} in $\tilde{f}...$

(small μ dependence)

Building a BFKL MC

... and rewrite the BFKL eq. as

$$(\omega - \omega_0) \tilde{f}(\vec{q}_{a\perp}, \vec{q}_{b\perp}, \omega) = \frac{1}{2} \delta(\vec{q}_{a\perp} - \vec{q}_{b\perp}) + \frac{\bar{\alpha}_s}{\pi} \int_{k_{\perp}^2 > \mu^2} \frac{d^2 \vec{k}_{\perp}}{k_{\perp}^2} \tilde{f}(\vec{q}_{a\perp} + \vec{k}_{\perp}, \vec{q}_{b\perp}, \omega)$$

$$\omega_0 = \frac{\bar{\alpha}_s}{\pi} \int \frac{d^2 \vec{k}_{\perp}}{k_{\perp}^2} \left[\theta(\mu^2 - k_{\perp}^2) - \frac{q_{a\perp}^2}{\vec{k}_{\perp}^2 + (\vec{q}_{a\perp} + \vec{k}_{\perp})^2} \right] = \bar{\alpha}_s \ln \left(\frac{\mu^2}{q_{a\perp}^2} \right).$$

ω_0 describes the net effect from unresolved & virtual gluon emission, and we are left with an integral over **resolved** gluons.

Solve by iteration and perform the inverse Mellin transform to get...

Building a BFKL MC

$$f(\vec{q}_{a\perp}, \vec{q}_{b\perp}, \Delta y) = \sum_{n=0}^{\infty} f^{(n)}(\vec{q}_{a\perp}, \vec{q}_{b\perp}, \Delta y)$$

$$f^{(0)}(\vec{q}_{a\perp}, \vec{q}_{b\perp}, \Delta y) = \left[\frac{\mu^2}{q_{a\perp}^2} \right]^{\bar{\alpha}_s \Delta y} \frac{1}{2} \delta(\vec{q}_{a\perp} - \vec{q}_{b\perp}),$$

$$f^{(n \geq 1)}(\vec{q}_{a\perp}, \vec{q}_{b\perp}, \Delta y)$$

$$= \left[\frac{\mu^2}{q_{a\perp}^2} \right]^{\bar{\alpha}_s \Delta y} \left\{ \prod_{i=1}^n \int d^2 \vec{k}_{i\perp} dy_i \mathcal{F}_i \right\} \frac{1}{2} \delta(\vec{q}_{a\perp} - \vec{q}_{b\perp} - \sum_{i=1}^n \vec{k}_{i\perp})$$


$$\mathcal{F}_i = \frac{\bar{\alpha}_s}{\pi k_{i\perp}^2} \theta(k_{i\perp}^2 - \mu^2) \theta(y_{i-1} - y_i) \left[\frac{(\vec{q}_{a\perp} + \sum_{j=1}^{i-1} \vec{k}_{j\perp})^2}{(\vec{q}_{a\perp} + \sum_{j=1}^i \vec{k}_{j\perp})^2} \right]^{\bar{\alpha}_s y_i}$$

BFKL eq. recast in terms of integrals over **resolved** gluon phase space.


Form factors describes the unresolved radiation. The full sum is still

μ -independent. Suitable for MC integration!

Does it work?




Hadron Cross Sections




$W + 2\text{-jet}$

New setup:




- New Impact Factor
- Same BFKL chain
- Better signal

But now the cross section depends on the quark **and** the gluon pdf. The quark pdf is less steep \rightarrow less pdf suppression \rightarrow BFKL *enhancement*




$W + 2\text{-jet}$



$$\Delta y = y_{j_2} - y_{j_1}, y_W, y_{j_2} \geq 1, y_{j_1} \leq -1$$

4b production (in progress)



Summary

The Monte Carlo method to study BFKL physics works!

1. It can reproduce (Naive) BFKL
2. It respects energy & momentum conservation
3. It can even include subleading logs!
4. It proved very efficient in incorporating experimental cuts to the momenta of the scattered particles...
5. ... and allows for a further study of the radiation from the BFKL ladder