



Low-energy enhancement in the γ -ray strength functions of $^{73,74}\text{Ge}$



Therese Renstrøm
Oslo Cyclotron Laboratory,
University of Oslo





Many people measuring the same property...?



Updated version of famous graph,
courtesy of Ann-Cecilie Larsen

Many people measuring the same property...?

H. Utsunomiya, Phys. Rev. C 88,015805 (2013)




P. E. Koehler, Phys. Rev. C 88, 041305(R) (2013)

-New neutron resonance parameters, small uncertainties



Need for a new common case?


^{74}Ge



$^{74}\text{Ge}(^3\text{He}, ^3\text{He}')$
@OCL



$^{74}\text{Ge}(p, p')$
@LLNL




$^{74}\text{Ge}(\gamma, \gamma')$
@ γELBE



$^{74}\text{Ge}(\alpha, \alpha')$
@iThemba


^{74}Ge



$^{74}\text{Ge}(^3\text{He}, ^3\text{He}')$
@OCL



$^{74}\text{Ge}(p, p')$
@LLNL



$^{74}\text{Ge}(\gamma, \gamma')$
@ γ ELBE



$^{74}\text{Ge}(\alpha, \alpha')$
@iThemba

^{74}Ge



The first step is compilation

1. Data sources

1. "Oslo method" ★
 2. (γ, γ') with Brehm ★
 3. (γ, γ') with HIGS
 4. Direct Reac.+ HPGe (DR-TSC) ★
 5. (n, γ) + statistical model
 6. $(n, 2\gamma)$
 7. (γ, n) – Utsonomya ★
 8. (p, p') , (α, α')
 9. (e, e')
 10. (n, n')
- Note: Many of these are inextricably linked with Level Density

2. Database format

3. Evaluation methodology

4. New measurements/Benchmark

5. Understanding sources of discrepancies



$^{74}\text{Ge}(p, p')$




$^{74}\text{Ge}(\gamma, \gamma')$





$^{74}\text{Ge}(\alpha, \alpha')$

Advantages by this type of joint effort:

- Timescale
- Communication
- Understanding of data analysis
- More FUN!





Why Germanium???





Level density and gamma-decay strength
can be extracted using the Oslo-method



Figure courtesy of Ann-Cecilie Larsen

“Almost all of the (n,γ) cross sections of the 277 stable isotopes have been measured. The few exceptions are ^{17}O , $^{36,38}\text{Ar}$, ^{40}K , ^{50}V , ^{70}Zn , $^{72,73}\text{Ge}$, $^{77,82}\text{Se}$, $^{98,99}\text{Ru}$, ^{131}Xe , ^{138}La , ^{158}Dy , and ^{195}Pt ...”, F. Käppeler et al., Rev. of Mod. Phys, 83, 2011






A. Spyrou, PRL **113**, 232502 (2014)





Two theoretical explanations of the type:

Argues for [E1](#)
E. Litvinova and N. Belov,
Phys. Rev. C 88, 031302(R)
(2013)

Argues for [M1](#)
R. Schwengner, S.
Frauendorf, and A. C.
Larsen, Phys. Rev.
Lett. 111, 23504 (2013)



Effect on (n, γ) reaction rates



A. C. Larsen and S. Goriely, Phys. Rev. C 82, 014318 (2010)


Two experiments below and above Sn



$(^3\text{He}, ^3\text{He}')$


$(^3\text{He}, \alpha)$

(γ, n)



^{74}Ge & bonus nucleus ^{73}Ge

E(NaI) : E(Si) strip 4




$(^3\text{He}, \alpha\gamma)$





$(^3\text{He}, ^3\text{He}\gamma)$




Ge-73



Ge-74





Neutron resonance data

-discrepancies between the libraries

Atlas of N. Res. by S.F. Mughabghab


RIPL3

In the case of $^{73,74}\text{Ge}$, s-wave spacings, **D0**, are given in both **Atlas of N. Res. by S.F. Mughabghab** and RIPL. After careful consideration we have chosen to use an **average value** of the two proposed sets of **D0** values and uncertainties. The two main reasons for this choice are:

1. For $^{73,74}\text{Ge}$ the **D0** values from **Atlas of N. Res.** are larger by 38% and 60%, respectively, than the values given in RIPL3.
2. **Atlas of N. Res.** presents a table of measured resonances giving a good transparency.

The experimental results that give the values listed in RIPL-3 are not presented in any peer-reviewed publication.


The photo-neutron experiment @NewSUBARU synchrotron radiation facility



Relativistic electrons on eV photons

Laser Compton backscattered
-ray beams are ideal because:

- Almost monochromatic
- Tunable energies




$$\gamma = E_e/mc^2$$

$$E_\gamma = \frac{4\gamma^2 \epsilon_L}{1 + (\gamma\theta)^2 + 4\gamma\epsilon_L/(mc^2)}$$







^{20}He proportional
counters




$$\int_{S_n}^{E_{\max}} n_{\gamma}(E_{\gamma}) \sigma(E_{\gamma}) dE_{\gamma} = \frac{N_n}{N_t N_{\gamma} \xi \epsilon_n g}$$






^{20}He proportional
counters




$$\int_{S_n}^{E_{\max}} n_{\gamma}(E_{\gamma})\sigma(E_{\gamma})dE_{\gamma} = \frac{N_n}{N_t N_{\gamma} \xi \epsilon_n g}$$




Solve integral using
Taylor expansion method



^{20}He proportional
counters




$$\int_{S_n}^{E_{\max}} n_{\gamma}(E_{\gamma})\sigma(E_{\gamma})dE_{\gamma} = \frac{N_n}{N_t N_{\gamma} \xi \epsilon_n g}$$





Solve integral using
Taylor expansion method



Photoneutron cross sections, ^{74}Ge



$$f \uparrow (E_\gamma) = \frac{1}{3\pi^2 \hbar^2 c^2} \frac{\sigma(E_\gamma)}{E_\gamma}$$

Detailed balance $f \uparrow = f \downarrow$
(allows us to put it all together)







Talk of M. Wiedeking ☺

THE LOW ENERGY
ENHANCEMENTS...


β -Oslo upbend



A. Spyrou, PRL **113**, 232502 (2014)





Shell-model calculations, ^{74}Ge





Talk of R. Schwengner


Level densities of $^{73,74}\text{Ge}$



S. Goriely et al., Phys. Rev. C 78 064307, (2008)





Constraining (n, γ)-cross sections





*TALYS: Koning et al., <http://www.talys.eu>

Using a variety of different inputs in TALYS




Maxwellian Average Cross Sections

Reaction	kT (keV)	MACS (mb) (new)	MACS(mb)*
$^{72}\text{Ge}(n,\gamma)$	30	66 (13)	73 (7)
$^{73}\text{Ge}(n,\gamma)$	30	294 (78)	243 (47)



The search for the upbend continues...



Beta-Oslo method on $^{78,80}\text{Ge}$


NSCL PAC 39 PROPOSAL

“Search for the upbend in neutron-rich Ge isotopes”




Spokespersons:

A. C. Larsen, M. Guttormsen, T. Renstrøm (University of Oslo, Norway)

A sneak peek at the most neutron rich isotope analysed to date using the beta-Oslo method...




Experiments performed in Oslo,
 γ SF to be published.



The β -Oslo method for ^{70}Ni

Discretionary beam time @ NSCL/MSU,
February 2015;

^{70}Co beta-decaying into ^{70}Ni



[S.N. Liddick et al., in preparation]

Summary

- Strength functions of $^{73,74}\text{Ge}$ both show a low energy enhancement
- If this enhancement is there for more neutron-rich Ge-isotopes, it will strongly influence the (n, γ) reaction rates
- Shell-model calculations indicate that the upbend is of M1-type
- Good agreement between the photo-neutron data and the Oslo data

THANK YOU FOR
LISTENING